

Prevalence of lung function impairment among Greek cement production workers: a cross-sectional study

George RACHIOTIS^{1*}, Konstantinos KOSTIKAS¹, Dimitra PINOTSI²,
Christos HADJICHRISTODOULOU¹ and Spyros DRIVAS²

¹Department of Hygiene and Epidemiology, Medical Faculty, University of Thessaly, Greece

²Greek Institute for Occupational Safety and Health, Greece

Received January 6, 2017 and accepted August 15, 2017

Published online in J-STAGE August 22, 2017

Abstract: Greece is a significant cement producing country. The aim of this study was to investigate the prevalence and risk factors of lung function impairment among Greek cement workers. One hundred thirty-seven cement production workers participated in this study. In addition, 110 employees not exposed to cement dust comprised the control group. The concentration of cement total dust at workplace varied from 1.1 to 11.6 mg/m³. In only one of the measurements, the Threshold Limit Level of 10 mg/m³ has been exceeded. Cement production workers presented a higher prevalence of FEV1 < 80% in comparison to controls (13.9% vs. 2.7%; Chi-Square Test; $p=0.002$). Multivariate analysis has shown that cement production workers have recorded an almost 5 fold risk of low lung function, as expressed by FEV1 < 80%, in comparison to the reference population (OR=4.92; 95% C.I.=1.22–12.62). Current smoking was associated with an almost 4-fold increased risk of FEV1 < 80% (OR=3.91; 95% C.I.=1.32–11.56). In conclusion, we found a high prevalence of impaired lung function among Greek cement production workers, despite the fact that total and inhalable dust levels were below the occupational exposure limits.

Key words: Cement, Dust, Worker, Spirometry, Greece

The production of cement is related to a process that includes multiple stages: quarrying, crushing, raw milling, blending, and production of clinker, milling and packing. It is well known that cement dust contains respirable particles with aerodynamic diameter ranged from 0.05 to 5 μm ¹. There is conflicting evidence on the association between exposure to cement dust and lung function impairment^{1,2}. Greece is a significant cement producing country³. To the best of our knowledge, there is no published information on the impact of the exposure to cement dust on the respiratory system in Greece. Consequently, the aim of our study was to investigate the prevalence and risk factors of lung function impairment among Greek cement workers.

Study population

A convenient sample (n=150) of blue collar cement workers at a Greek cement production plant was invited to participate in the cross-sectional study. One hundred thirty-seven cement production workers participated in the study (response rate: 91%). A convenient sample of 110 employees not exposed to cement dust comprised the reference (control) group. The control group consisted of office (white collar) employees outside the cement industry plant. The members of this group performed various administrative and clerical tasks.

Environmental monitoring

The assessment of the environmental exposure to cement dust (total and respirable fraction) for the study group was performed by the use of the gravimetric method. In particular, the air was drawn through a cellulose filter of known weight that was adapted in a personal sampler device that

*To whom correspondence should be addressed.

E-mail: g.rachiotis@gmail.com

©2018 National Institute of Occupational Safety and Health

was attached to a sampling pump of fixed flow (MSA Company, Pittsburgh, Pennsylvania, USA). Reweighing of the filter, after sampling, gives a measurement of the dust weight, and thus of dust concentration at the workplace. In addition, a cyclone was used to separate respirable and oversize dust. Total or inhalable dust refers to dust particles with an aerodynamic diameter of 100 μm . Respirable dust refers to particles with an aerodynamic diameter of 5 μm as well as smaller particles. In total, 14 measurements of total and 22 measurements of respirable dust were performed.

Spirometry

All participants underwent spirometry by using a portable spirometer (Vitalograph, Buckingham, England). Spirometries were performed according to the ERS guidelines⁴). Three technically satisfactory maximal forced exhalations were recorded and the best forced expiratory volume in one second (FEV1), forced vital capacity (FVC), and the ratio FEV1/FVC were taken into account for statistical analysis. Predictive values of the above parameters were extracted based on the recommendations of the European Coal and Steel Community. The evaluation of extracted values regarding the parameters of spirometry was based on GOLD criteria⁵). In addition, the portable spirometer was calibrated with a three-l syringe.

Statistical analysis

Continuous variables are presented as mean (SD) and categorical variables as absolute (n) and relative (%) frequencies. Associations between categorical variables were explored by the use of χ^2 test (univariate analysis). Continuous variables were tested for normality by Kolmogorov-Smirnov test. Univariate analysis of normally distributed continuous variables was performed by student's *t*-test. Logistic regression analysis was used as the multivariate analysis in order to assess the impact on symptomatic status, and respiratory function of occupational exposure to cement dust, smoking use, and age. In these models of multivariate analysis respiratory symptoms and FEV1 < 80%, were the dependent variable, while age, smoking, and occupational exposure to cement dust, were the independent ones. Odds ratios (OR) and 95% confidence interval (95% CI) were calculated. The level of statistical significance was set at 0.05. All statistical tests were two sided. Statistical analyses were performed with Epi info software.

Ethics

The protocol of the study has been approved by the Steering Committee of the Post Graduate Program

“Applied Public and Environmental Hygiene” at the University of Thessaly, Greece.

The concentration of cement total dust at workplace varied from 1.1 to 11.6 mg/m^3 . In only one of the measurements the Threshold Limit Level established in Greece of 10 mg/m^3 has been exceeded. The concentration of the respirable fraction of cement dust was below the national Threshold Limit Value of 5 mg/m^3 , and ranged between 0.1 to 3.4 mg/m^3 . The vast majority of the participants ($n=198$; 80%) were males and 49 (20%) females. The mean age of our sample was 47 yr old (SD=19) and the mean of duration of employment was 19 yr (SD=9.4). Regarding occupational exposure status, 137 (55.5%) participants were cement production workers and 110 (45.5%) comprised the control group. The gender and age distribution was similar in both exposure and control group. The overall prevalence of current smoking among participants was estimated at 56% (54.7% among cement production workers and 58.6% among controls; $p=\text{ns}$). Cement workers reported a significantly higher prevalence of respiratory symptoms in comparison to office workers (20% vs. 4%, respectively, p value < 0.05). Table 1 depicts the mean values of FEV1, FVC and FEV1/FVC by age group, smoking, occupational exposure status and duration of employment. The prevalence of the patients with a ratio FEV1/FVC < 70% was numerically higher in cement vs. office workers (4.4% vs. 1.1%, respectively); however this difference was not statistically significant ($p=0.1$). Cement workers additionally demonstrated higher FVC mean value than the control group. Cement production workers, participants with age < 47 yr old, and current smokers have recorded in significantly lower mean values of FEV1. The univariate analysis of FEV1 < 80% (Table 2) has shown that cement production workers and current smokers had a higher risk to present an FEV1 < 80%. In particular, the homogeneous group of cement production workers presented a higher prevalence of FEV1 < 80% in comparison to the subsample of controls (13.9% vs. 2.7%; χ^2 Test; $p=0.002$). The prevalence of FEV1 < 80% increased with duration of employment and also with age. Current smokers have had a higher prevalence of FEV1 < 80% in comparison to non-current smokers (12.2% vs. 4.6%, respectively; χ^2 Test; $p=0.038$). Table 3 presents the results of the logistic regression analysis of FEV1 < 80%. Cement production workers have recorded an almost 5 fold risk of FEV1 < 80% in comparison to the reference population OR = 4.92; 95% C.I. = 1.22 – 12.62). Current smoking was associated with an almost 4- fold increased risk of FEV1 < 80% (OR = 3.91; 95% C.I. = 1.32 – 11.56).

The present cross-sectional workplace-based study sug-

Table 1. Distribution of selected spirometric mean values in relation to participants' socio-demographic features (n=247)

Socio-demographic features	SPIROMETRIC VALUES					
	FEV1/FVC (% predicted)		FEV1 (% predicted)		FVC (% predicted)	
	Mean	SD	Mean	SD	Mean	SD
Smoking						
Current smokers	96.4	6.7	97.1	16.2	100.8	16
Non current smokers	97.2	8.9	99.3	15.5	100.9	15.2
Occupational status						
Exposed to cement dust	96.1	8.8	94.3	15.4	100.4	11.0
Non exposed to cement dust	97.6	6.0	95.8	15.8	99.4	15.0
Age						
≤47 yr	97.3	5.6	95.0	15.0	102	14.5
>47 yr	95.8	10.1	94.0	16.0	99.4	17.2
Duration of employment						
≤ 19 yr	97.2	5.5	99.2	12.8	101.4	13.4
> 19 yr	96.3	13.2	96.9	18.3	100.3	17.6

Table 2. Univariate analysis of FEV1<80%

Socio-demographic features	PREVALENCE	
	FEV1 (% predicted) < 80	≥80
Smoking		
Current smokers	12.2%*	87.8%
Non-current smokers	4.6%	95.4%
Occupational status		
Exposed to cement	13.9%*	86.1%
Non exposed to cement	2.7%	97.3%
Age		
≤47 yr	5.4%*	94.6%
>47 yr	14.1%	85.9%
Duration of employment		
≤ 19 yr	5.0%	95.0%
> 19 yr	12.6%	87.4%

**p*<0.05

gests that cement production workers have recorded elevated prevalence of abnormal lung function in comparison to the control population. The evidence on the association between exposure to cement dust and lung function impairment is controversial. There are several studies suggested a positive association⁶⁻¹² whereas other found no association^{13, 14}. However, it is of note that two recent well-conducted multi-national studies found a strong association between exposure to cement dust and lung function decline. In particular, Nordby *et al.*, in a multinational study supported by the European Cement Association (CEMBUREAU) reported reduced lung volumes among cement dust exposed workers¹⁵. In this context, a prospective study of workers employed in 24 cement production plants from 8 countries found that occupational exposure to cement dust was associated with reduced lung function parameters¹⁶. It is worth mentioning that we observed increased lung func-

Table 3. Multivariate analysis of FEV1<80%

Variable	FEV1/80%		<i>p</i>
	OR	95% CI	
Smoking			
Current smokers	3.91	1.32–11.56	0.014
Non-current smokers	1.00 (ref)		
Occupational status			
Exposed to cement	4.91	1.22–12.62	0.025
Non exposed to cement	1.00 (ref)		
Age			
>47 yr	3.19	0.79–12.88	NS
≤47 yr	1.00 (ref)		
Duration of employment			
> 19 yr	1.00 (ref)		NS
≤ 19 yr	0.69	0.14–3.30	NS

NS: non significant

tion impairment among cement production workers despite the fact that the measured levels of total and respirable dust were below the national occupational exposure limits. It has been suggested by other researchers that the current occupational exposure limit for total dust (10 mg/m³) is too high and should be reduced to protect the respiratory health of cement production workers⁶. We also found a high prevalence (56%) of current smoking among blue collar cement production workers. This finding corroborates previous studies among heavy industry workers in Greece^{17, 18}. Since smoking is a significant contributing factor of lung function impairment, anti-smoking activities targeting blue collar workers may be warranted. Our study presents several limitations which need consideration. A first limitation is the cross-sectional design of the survey. A second limitation is related to sampling method used in our study. Our sample was convenient, and the results are hardly applicable across

the board of all cement workers in Greece. An additional potential limitation of our study could be related to “healthy worker effect”. In particular, ill cement production workers might have left exposed jobs, while healthy employees may have remained in jobs that could be hazardous. This selection bias could have resulted in an underestimation of the effect of cement dust on lung function parameters. Another limitation refers to the absence of longitudinal (historical) exposure data on the levels of cement dust. Last, despite the use of logistic regression analysis, we can’t exclude a residual confounding effect of occupational status (blue vs. white collar workers) on FEV1. In conclusion, we found a high prevalence of obstructive ventilation pattern among Greek cement production workers despite the fact that total and inhalable dust levels were below the occupational exposure limits. The significant association between occupational exposure to cement dust and obstructive lung function ventilation pattern was independent of the impact of smoking on lung function indices.

References

- 1) Meo SA (2004) Health hazards of cement dust. *Saudi Med J* **25**, 1153–9.
- 2) Ahmed HO, Abdullah AA (2012) Dust exposure and respiratory symptoms among cement factory workers in the United Arab Emirates. *Ind Health* **50**, 214–22.
- 3) Hellenic Cement Industry Association. <http://www.hcia.gr/el/statistical-elements/production/>
- 4) Miller MR, Hankinson J, Brusasco V, Burgos F, Casaburi R, Coates A, Crapo R, Enright P, van der Grinten CP, Gustafsson P, Jensen R, Johnson DC, MacIntyre N, McKay R, Navajas D, Pedersen OF, Pellegrino R, Viegi G, Wanger J; ATS/ERS Task Force (2005) Standardisation of spirometry. *Eur Respir J* **26**, 319–38.
- 5) Pauwels RA, Buist AS, Calverley PM, Jenkins CR, Hurd SS; GOLD Scientific Committee (2001) Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease. NHLBI/WHO Global Initiative for Chronic Obstructive Lung Disease (GOLD) Workshop summary. *Am J Respir Crit Care Med* **163**, 1256–76.
- 6) Mwaiselage J, Bråtveit M, Moen B, Mashalla Y (2004) Cement dust exposure and ventilatory function impairment: an exposure-response study. *J Occup Environ Med* **46**, 658–67.
- 7) Oleru UG (1984) Pulmonary function and symptoms of Nigerian workers exposed to cement dust. *Environ Res* **33**, 379–85.
- 8) Noor H, Yap CL, Zolkepli O, Faridah M (2000) Effect of exposure to dust on lung function of cement factory workers. *Med J Malaysia* **55**, 51–7.
- 9) Meo SA, Al-Drees AM, Al Masri AA, Al Rouq F, Azeem MA (2013) Effect of duration of exposure to cement dust on respiratory function of non-smoking cement mill workers. *Int J Environ Res Public Health* **10**, 390–8.
- 10) Yang CY, Huang CC, Chiu HF, Chiu JF, Lan SJ, Ko YC (1996) Effects of occupational dust exposure on the respiratory health of Portland cement workers. *J Toxicol Environ Health* **49**, 581–8.
- 11) Zeleke ZK, Moen BE, Bråtveit M (2010) Cement dust exposure and acute lung function: a cross shift study. *BMC Pulm Med* **10**, 19.
- 12) Zeleke ZK, Moen BE, Bråtveit M (2011) Lung function reduction and chronic respiratory symptoms among workers in the cement industry: a follow up study. *BMC Pulm Med* **11**, 50.
- 13) Abrons HL, Petersen MR, Sanderson WT, Engelberg AL, Harber P (1988) Symptoms, ventilatory function, and environmental exposures in Portland cement workers. *Br J Ind Med* **45**, 368–75.
- 14) Fell AK, Thomassen TR, Kristensen P, Egeland T, Kongerud J (2003) Respiratory symptoms and ventilatory function in workers exposed to portland cement dust. *J Occup Environ Med* **45**, 1008–14.
- 15) Nordby KC, Fell AK, Notø H, Eduard W, Skogstad M, Thomassen Y, Bergamaschi A, Kongerud J, Kjuus H (2011) Exposure to thoracic dust, airway symptoms and lung function in cement production workers. *Eur Respir J* **38**, 1278–86.
- 16) Nordby KC, Notø H, Eduard W, Skogstad M, Fell AK, Thomassen Y, Skare Ø, Bergamaschi A, Pietroiusti A, Abderhalden R, Kongerud J, Kjuus H (2016) Thoracic dust exposure is associated with lung function decline in cement production workers. *Eur Respir J* **48**, 331–9.
- 17) Rachiotis G, Behrakis PK, Vasiliou M, Yfantopoulos J (2006) Quality of life and smoking among industrial workers in Greece. *Med Lav* **97**, 44–50.
- 18) Drivas S, Rachiotis G, Vlastos FD, Zacharias C, Alexopoulos CG, Symvoulakis M, Vasiliou M, Behrakis PK (2007) Occupational exposure to lignite and impact on respiratory system among heavy industry personnel. *Ind Health* **45**, 409–14.